

A SUPPORT THEOREM FOR THE RADIATION FIELDS ON ASYMPTOTICALLY EUCLIDEAN MANIFOLDS

ANTÔNIO SÁ BARRETO

1. INTRODUCTION

We prove a support theorem for the radiation fields on asymptotically Euclidean manifolds with metrics which are warped products near infinity. It generalizes to this setting the well known support theorem for the Radon transform in \mathbb{R}^n . The main reason we are interested in proving such a theorem is the possible application to the problem of reconstructing an asymptotically Euclidean manifold from the scattering matrix at all energies, see [14].

An asymptotically Euclidean manifold [12, 8] is a C^∞ compact manifold X with boundary ∂X , which is equipped with a C^∞ Riemannian metric g that in a collar neighborhood of the boundary ∂X satisfies

$$(1.1) \quad g = \frac{dx^2}{x^4} + \frac{h(x)}{x^2}, \quad \text{in } [0, \epsilon) \times \partial X,$$

where x is a defining function of ∂X and h is a C^∞ one parameter family of metrics on ∂X . The basic example is the radial compactification of \mathbb{R}^n , [12].

In this paper we will consider the class of metrics g which have the following special form near ∂X :

$$(1.2) \quad g = \frac{dx^2}{x^4} + \psi(x) \frac{h_0}{x^2}, \quad x \in [0, \epsilon),$$

where $\psi \in C^\infty([0, \epsilon))$, $\psi(x) > 0$, $\psi(0) = 1$, and h_0 is a C^∞ metric on ∂X . These are known as warped product metrics [11].

We consider the wave equation on X . Let Δ_g be the Laplace operator on X , and let $u(t, z)$ satisfy

$$(1.3) \quad \begin{aligned} (D_t^2 - \Delta_g)u(t, z) &= 0 \text{ on } \mathbb{R} \times X, \\ u(0) &= f_1, \quad D_t u(0) = f_2, \quad \text{with } f_1, f_2 \in C_0^\infty(X). \end{aligned}$$

A function $f \in C_0^\infty(X)$ if it is C^∞ and its support does not intersect the boundary of X .

The following is proved in [4, 5]:

Theorem 1.1. *Let x be the boundary defining function for which (1.1) holds, and let $z = (x, y)$, $y \in \partial X$, be the corresponding boundary normal coordinates in a collar neighborhood of the boundary. Then*

$$(1.4) \quad \begin{aligned} v_+(x, s, y) &= x^{-\frac{n-1}{2}} u(s + \frac{1}{x}, x, y) \in C^\infty(\mathbb{R}_s \times [0, \epsilon)_x \times \partial X), \\ v_-(x, s, y) &= x^{-\frac{n-1}{2}} u(s - \frac{1}{x}, x, y) \in C^\infty(\mathbb{R}_s \times [0, \epsilon)_x \times \partial X). \end{aligned}$$

Friedlander [4, 5] defined the forward and backward radiation fields respectively as

$$(1.5) \quad \mathcal{R}_+(f_1, f_2) = D_s v_+(0, s, y) \quad \text{and} \quad \mathcal{R}_-(f_1, f_2) = D_s v_-(0, s, y).$$

Lax and Phillips [9] proved that in \mathbb{R}^n the forward (or backward) radiation field is the modified Radon transform, that is:

$$\mathcal{R}_+(f_1, f_2)(s, \omega) = |D_s|^{\frac{n-3}{2}} Rf_1(s, \omega) + |D_s|^{\frac{n-1}{2}} Rf_2(s, \omega), \text{ where}$$

$$Rf(s, \omega) = \int_{\langle x, \omega \rangle = s} f(z) d\sigma, \quad \sigma \text{ is the surface measure on } \langle x, \omega \rangle = s, \text{ is the Radon transform.}$$

Helgason's celebrated support theorem for Radon transforms [6] says that if f is a rapidly decaying function in \mathbb{R}^n and $Rf(s, \omega) = 0$ for $s < s_0$, $s_0 < 0$ (and hence by symmetry $Rf(s, \omega) = 0$ for $|s| > |s_0|$) then f is supported in the ball of radius $|s_0|$. The assumption that f is rapidly decaying cannot be entirely removed. For example, for any $m \in \mathbb{N}$, there are smooth functions $f(z)$ in $z \in \mathbb{R}^n$ which are not compactly supported, decay like $|z|^{-m}$, and whose Radon transform is compactly supported. See for example [6].

This is a result in control theory, where the support of a function can be exactly controlled by the support of its Radon transform. We want to address the analogue question for Radiation fields. The following was proved in [13]:

Theorem 1.2. *If $f \in C_0^\infty(X)$, g is an arbitrary asymptotically Euclidean metric, and $\mathcal{R}(0, f)(s, y) = 0$ for $s < -\frac{1}{x_0}$, $x_0 \in (0, \epsilon)$, then $f = 0$ if $x < x_0$.*

This says that if there exists some $x_1 \in (0, \epsilon)$, such that $f(x, y) = 0$ for $x < x_1$, but $\mathcal{R}_+(0, f)(s, y) = 0$ for $s < -\frac{1}{x_0}$, and $x_0 > x_1$, then in fact $f(x, y) = 0$ if $x < x_0$. The purpose of this paper is to discuss the following

Question 1.1. *Let (X, g) be an asymptotically Euclidean manifold, and let $\mathcal{S}(X)$ be the space of functions in $C^\infty(X)$ which are smooth up to ∂X and vanish to infinite order at ∂X . If $f \in \mathcal{S}(X)$ and $\mathcal{R}_+(0, f)(s, y) = 0$ for $s < -\frac{1}{x_0}$, $x_0 \in (0, \epsilon)$, is it true that $f = 0$ if $x < x_0$?*

We answer this question in the affirmative in the following particular case:

Theorem 1.3. *If g is a warped product metric, the dimension of X is greater than or equal to 3, $f \in \mathcal{S}(X)$, and $\mathcal{R}_+(0, f)(s, y) = 0$ for $s < -\frac{1}{x_0}$, then $f = 0$ if $x < x_0$.*

2. ENERGY ESTIMATES

The first step in the proof of Theorem 1.3 is to obtain estimates the solution to (1.3) up to $x = 0$ and $s = -\infty$.

The Laplacian with respect to the metric (1.2) is, in a neighborhood of ∂X , given by

$$\Delta_g = -x^4 \partial_x^2 + (n-3)x^3 \partial_x - x^4 A(x) \partial_x + x^2 \psi(x)^{-1} \Delta_{h_0},$$

where $A(x) = \partial_x \log \left(\psi(x)^{\frac{n-1}{2}} \right)$, and Δ_{h_0} is the Laplacian on ∂X with respect to the metric h_0 . In what follows it is convenient to get rid of first order terms, so we will work with $Q = F(x)^{-1} \Delta_g F(x)$, with $F(x) = x^{\frac{n-1}{2}} (\psi(x))^{-\frac{n-1}{4}}$. We get that

$$Q = -(x^2 \partial_x)^2 + x^2 \phi(x) \Delta_{h_0} + x^2 B(x), \text{ where}$$

$$\phi(x) = [\psi(x)]^{-1}, \quad B(x) = \frac{(n-1)(n-3)}{4} + x B_1(x), \quad B_1 \in C^\infty([0, \epsilon)).$$

The wave equation (1.3) is translated into

$$(2.1) \quad (\partial_t^2 - (x^2 \partial_x)^2 + x^2 \phi(x) \Delta_{h_0} + x^2 B(x))u = 0,$$

$$u(0) = f_1(x, y), \quad \partial_t u(0) = f_2(x, y).$$

Instead of working with coordinates x and s , and the forward and backward radiation fields separately, it is better to work with the forward and backward radiation fields simultaneously. So we define

$$(2.2) \quad s_+ = t - \frac{1}{x} \text{ and } s_- = t + \frac{1}{x}, \quad x > 0.$$

Let u be the solution to the wave equation (2.1), and let $w = F(x)u$. In coordinates (2.2),

$$w(s_+, s_-, y) = F\left(\frac{2}{s_- - s_+}\right)u\left(\frac{s_+ + s_-}{2}, \frac{2}{s_- - s_+}, y\right)$$

satisfies

$$(2.3) \quad \begin{aligned} (s_- - s_+)^2 \partial_{s_+} \partial_{s_-} w + \phi\left(\frac{2}{s_- - s_+}\right) \Delta_{h_0} w + B\left(\frac{2}{s_- - s_+}\right) w &= 0, \\ w(s_+, -s_+, y) &= F\left(-\frac{1}{s_+}, y\right) f_1\left(-\frac{1}{s_+}, y\right), \\ (\partial_{s_+} w)(s_+, -s_+, y) &= \frac{1}{2} F\left(-\frac{1}{s_+}\right) f_2\left(-\frac{1}{s_+}, y\right) + \frac{1}{2s_+^2} F'\left(-\frac{1}{s_+}\right) f_1\left(-\frac{1}{s_+}, y\right) + \frac{1}{2s_+^2} F\left(-\frac{1}{s_+}\right) f_1'\left(-\frac{1}{s_+}, y\right), \end{aligned}$$

We want to understand the behavior of w as $s_+ \sim -\infty$, and $s_- \sim \infty$ and thus we compactify $\mathbb{R} \times X$ by setting

$$\mu = -\frac{1}{s_+} \text{ and } \nu = \frac{1}{s_-}$$

So

$$(2.4) \quad w(\mu, \nu, y) = F\left(\frac{2\mu\nu}{\mu + \nu}\right) u\left(\frac{\mu - \nu}{\mu + \nu}, \frac{2\mu\nu}{\mu + \nu}, y\right)$$

satisfies

$$(2.5) \quad \begin{aligned} \left((\mu + \nu)^2 \partial_\mu \partial_\nu - \phi\left(\frac{2\mu\nu}{\mu + \nu}\right) \Delta_{h_0} w - B\left(\frac{2\mu\nu}{\mu + \nu}\right) \right) w &= 0, \\ w(\mu, \mu, y) &= \tilde{f}_1(\mu, y), \quad (\partial_\mu w)(\mu, \mu, y) = \tilde{f}_2(\mu, y), \end{aligned}$$

where

$$\begin{aligned} \tilde{f}_1(\mu, y) &= F(\mu) f_1(\mu, y), \text{ and} \\ \tilde{f}_2(\mu, y) &= \frac{1}{2\mu^2} F(\mu) f_2(\mu, y) + \frac{1}{2} F'(\mu) f_1(\mu, y) + \frac{1}{2} F(\mu) f_1'(\mu, y). \end{aligned}$$

Since we are dealing with a degenerate equation, we will work with weighted Sobolev spaces.

Definition 2.1. Let $T > 0$ and $\Omega_T = (0, T) \times (0, T)$. We define

$$\begin{aligned} H_j^s([0, T] \times \partial X) &= \{f \in L^2([0, T] \times \partial X), \quad \mu^{-j} f \in H^s([0, T] \times \partial X)\}, \text{ with norm } \|f\|_{s,j}, \text{ and} \\ H_j^s(\Omega_T \times \partial X) &= \{f \in L^2(\Omega_T \times \partial X), \quad (\mu + \nu)^{-j} f \in H^s(\Omega_T \times \partial X)\}, \text{ with norm } \|f\|_{s,j}, \text{ and} \end{aligned}$$

The next step is to prove

Theorem 2.2. Let $\tilde{f}_j(\mu, y) \in C^\infty([0, T] \times \partial X)$, $j = 1, 2$, be such that $\partial_\mu^k \tilde{f}_j(0, y) = 0$, $k = 0, 1, 2, \dots$. Let w satisfy (2.5) in $\Omega_T \times \partial X$. Then there exists $T_0 > 0$ such that w has a C^∞ extension up to $\overline{\Omega_T} \times \partial X$, $T \leq T_0$.

Proof. By finite speed of propagation, $w \in C^\infty(\Omega_T \times \partial X)$, we want to establish the regularity up to the closure $\overline{\Omega_T} \times \partial X$. We begin the proof with the following elementary lemmas:

Lemma 2.3. *Let $\mu \in (0, T)$, and $w \in C^\infty(\Omega_T)$. Then the following inequalities are true*

$$(2.6) \quad \int_{\mu}^b (\mu + \nu)^{-1-k} |w(\mu, \nu)|^2 d\nu \leq 2b\mu^{-1-k} |w(\mu, \mu)|^2 + b \int_{\mu}^b (\mu + \nu)^{-k} |\partial_{\nu} w(\mu, \nu)|^2 d\nu, \quad k \in \mathbb{N},$$

and

$$(2.7) \quad \int_{\mu}^{\nu} (\mu + \nu)^{-1} |w(\mu, \nu)|^2 d\mu \leq 2\nu^{-1} |w(\nu, \nu)|^2 + \nu \int_a^{\nu} |\partial_{\mu} w(\mu, \nu)|^2 d\mu, \quad k \in \mathbb{N}.$$

Proof. We write

$$(2.8) \quad w(\mu, \nu) = w(\mu, \mu) + \int_{\mu}^{\nu} \partial_s w(\mu, s) ds.$$

The Cauchy-Schwartz inequality gives

$$|w(\mu, \nu)|^2 \leq 2|w(\mu, \mu)|^2 + 2(\nu - \mu) \int_{\mu}^{\nu} |\partial_s w(\mu, s)|^2 ds.$$

Since $\nu > 0$

$$\begin{aligned} \int_{\mu}^b (\mu + \nu)^{-1-k} |w(\mu, \nu)|^2 d\nu &\leq 2(b - \mu)\mu^{-1-k} |w(\mu, \mu)|^2 + 2 \int_{\mu}^b \int_{\mu}^{\nu} (\mu + \nu)^{-k} |\partial_s w(\mu, s)|^2 ds d\nu \leq \\ &2b\mu^{-1-k} |w(\mu, \mu)|^2 + b \int_{\mu}^b (\mu + \nu)^{-k} |\partial_s w(\mu, s)|^2 d\nu. \end{aligned}$$

The proof of (2.7) is identical, and this ends the proof of the Lemma. \square

Lemma 2.4. *Let $\Omega_{a,b} \subset \Omega_T$ be the region defined by*

$$(2.9) \quad \Omega_{a,b} = \{(\mu, \nu) : \nu \geq \mu \geq 0, \quad 0 \leq a \leq \mu, \quad \nu \leq b\},$$

let $F(\mu, \nu) \in L^1(\Omega_T)$ and let $a_0 \leq a \leq b$. Then

$$(2.10) \quad \int_{a_0}^b \left(\int_{\Omega_{ab}} F(\mu, \nu) d\mu d\nu \right) da = \int_{\Omega_{a_0b}} (\mu - a_0) F(\mu, \nu) d\mu d\nu.$$

The proof is a straightforward application of Fubini's theorem.

Lemma 2.5. *If $\int_0^T \mu^{-m-2} |w(\mu, \mu)|^2 d\mu$ and $\int_{\Omega_{0T}} (\mu + \nu)^{-m} |(\partial_{\nu} - \partial_{\mu})w|^2 d\mu d\nu$ are finite, then*

$$(2.11) \quad \begin{aligned} &\int_0^T \int_0^T (\mu + \nu)^{-m-2} |w|^2 d\mu d\nu \leq \\ &2^{-m} \int_0^T \mu^{-m-1} |w(\mu, \mu)|^2 d\mu + \int_0^T \int_0^T (\mu + \nu)^{-m} |(\partial_{\nu} - \partial_{\mu})w|^2 d\mu d\nu. \end{aligned}$$

Proof. To see this it is better to rotate the axes μ and ν and use coordinates

$$(2.12) \quad \begin{aligned} r &= \mu + \nu, \quad \tau = \nu - \mu, \quad \text{so} \\ \mu &= \frac{1}{2}(r - \tau), \quad \nu = \frac{1}{2}(r + \tau). \end{aligned}$$

In the region $\mu \geq 0$, and $T \geq \nu \geq \mu$, we have

$$T\sqrt{2} \geq r \geq \tau, \quad \tau \geq 0.$$

In these coordinates $\partial_{\nu} - \partial_{\mu} = 2\partial_{\tau}$ and the diagonal $\mu = \nu$ becomes $\tau = 0$. We write, for $r \geq \tau$,

$$w(\tau, r) = w(0, r) + \int_0^{\tau} \partial_s w(s, r) ds$$

Using the Cauchy-Schwartz inequality and the fact that $\tau/r \leq 1$, we have

$$\begin{aligned} r^{-m-2}|w(\tau, r)|^2 &\leq 2r^{-m-2}|w_k(0, r)|^2 + 2\tau r^{-m-2} \int_0^\tau |\partial_s w_k(s, r)|^2 ds \leq \\ &2r^{-m-2}|w_k(0, r)|^2 + 2r^{-m-1} \int_0^\tau |\partial_s w_k(s, r)|^2 ds \end{aligned}$$

Therefore

$$\begin{aligned} \int_0^{T\sqrt{2}} \int_0^r r^{-m-2}|w^2(\tau, r)|^2 d\tau dr &\leq 2 \int_0^{T\sqrt{2}} r^{-m-1}|w(0, r)|^2 dr + 2 \int_0^{T\sqrt{2}} \int_0^r r^{-m-1} \int_0^\tau |\partial_s w_k(s, r)|^2 ds d\tau dr = \\ &2 \int_0^{T\sqrt{2}} r^{-m-2}|w(0, r)|^2 dr + 2 \int_0^{T\sqrt{2}} \int_0^r r^{-m-1}(r-s)|\partial_s w(s, r)|^2 ds dr. \end{aligned}$$

We get the same bound in the region $r \geq -\tau$. Translating this back into coordinates μ and ν we get (2.11). \square

Now we prove uniform energy estimates up to $\{\mu = 0\}$, $\{\nu = 0\}$. When $n > 3$ one can choose T_0 such that if $\mu < T_0$ and $\nu < T_0$, $B > \frac{(n-1)(n-3)}{8}$. When $n = 3$, B is not necessarily positive. In this case it is convenient to work with an eigenfunction decomposition. Let ϕ_k , $k \in \mathbb{N}$, be the eigenfunctions of Δ_{h_0} and let $\{\lambda_k\}$, $k \in \mathbb{N}$, with $0 = \lambda_{k-1} \leq \lambda_k$ be the corresponding eigenvalues. Let w be the solution to (2.5) and let $w_k = \langle w, \phi_k \rangle_{L^2(\partial X)}$. When $n = 3$ and $k = 1$, w_1 satisfies

$$(2.13) \quad \begin{aligned} \left((\mu + \nu)^2 \partial_\mu \partial_\nu - \frac{2\mu\nu}{\mu + \nu} B_1 \left(\frac{2\mu\nu}{\mu + \nu} \right) \right) w_1 &= 0 \text{ in } (0, T) \times (0, T), \\ w(\mu, \mu) &= q_1(\mu), \quad \partial_\mu w(\mu, \mu) = q_2(\mu), \end{aligned}$$

with $q_1 = \langle \tilde{f}_1, \phi_1 \rangle$ and $q_2 = \langle \tilde{f}_2, \phi_1 \rangle$. It is clear that $\|q_1\|_{s,j} < \infty$ and $\|q_2\|_{s,j} < \infty$ for every s and j .

We have

Proposition 2.6. *Let $\Omega_T = (0, T) \times (0, T)$ and let $w_1 \in C^\infty(\Omega_T)$ satisfy (2.13) in Ω_T . Suppose that $\|q_1\|_{s,j} < \infty$ and $\|q_2\|_{s,j} < \infty$, for all s and j . Then there exists $T_0 > 0$ such that $w_1 \in C^\infty(\overline{\Omega_T})$, $T \leq T_0$.*

Proof. The proof relies on the following energy estimates:

Lemma 2.7. *Let $w_1 \in C^\infty(\Omega_T)$ satisfy (2.13) in Ω_T . There exist T_0 and a constant $C = C(T_0) > 0$ such that if $T \leq T_0$,*

$$(2.14) \quad \begin{aligned} \text{for any fixed } \mu \in [0, T], \quad \int_0^T |\partial_\nu w_1(\mu, \nu)|^2 d\nu &\leq C(\|q_1\|_{1, \frac{1}{2}}^2 + \|q_2\|_{1, \frac{1}{2}}^2), \\ \text{and for any fixed } \nu \in [0, T], \quad \int_0^T |\partial_\mu w_1(\mu, \nu)|^2 d\mu &\leq C(\|q_1\|_{1, \frac{1}{2}}^2 + \|q_2\|_{1, \frac{1}{2}}^2). \end{aligned}$$

Proof. To prove this we first multiply (2.13) by $(\mu + \nu)^{-2}(\partial_\mu - \partial_\nu)w_1$. We obtain

$$\frac{1}{2} \partial_\nu ((\partial_\mu w_1)^2 + 2\mu\nu(\mu + \nu)^{-3} B_1 w_1^2) - \frac{1}{2} \partial_\mu ((\partial_\nu w_1)^2 + 2\mu\nu(\mu + \nu)^{-3} B_1 w_1^2) - \frac{1}{2} (\mu + \nu)^{-2} B_2 w_1^2 = 0,$$

where $B_2 = 2\frac{\mu-\nu}{\mu+\nu} B_1 + 4\frac{\mu\nu(\mu-\nu)}{(\mu+\nu)^2} B_1'(\frac{2\mu\nu}{\mu+\nu})$. After we integrate it in Ω_{ab} we obtain,

$$\begin{aligned} \frac{1}{2} \int_a^b ((\partial_\nu w_1)^2 + 2\mu\nu(\mu + \nu)^{-3} B_1 |w_1|^2) (a, \nu) d\nu + \frac{1}{2} \int_a^b ((\partial_\mu w_1)^2 + 2\mu\nu(\mu + \nu)^{-3} B_1 |w_1|^2) (\mu, b) d\mu - \\ \frac{1}{2} \int_{\Omega_{ab}} (\mu + \nu)^{-2} B_2 |w_1|^2 d\mu d\nu = \frac{1}{2\sqrt{2}} \int_a^b \left((q_1' + q_2)^2 + q_2^2 + \frac{1}{2} \mu^{-1} B_1(\mu) q_1^2 \right) d\mu. \end{aligned}$$

We apply Lemma 2.3 to show that we can pick T_0 such that for $T < T_0$,

$$\begin{aligned} \int_a^b 2\mu\nu(\mu+\nu)^{-3}|B_1||w_1|^2(a,\nu)d\nu &\leq 2a^{-1}q_1^2(a) + \frac{1}{4}\int_a^b (\partial_\nu w_1)^2(a,\nu)d\nu, \text{ and} \\ \int_a^b 2\mu\nu(\mu+\nu)^{-3}|B_1||w_1|^2(\mu,b)d\mu &\leq 2b^{-1}q_1^2(b) + \frac{1}{4}\int_a^b (\partial_\nu w_1)^2(\mu,b)d\mu. \end{aligned}$$

Therefore

$$(2.15) \quad \begin{aligned} &\frac{1}{4}\int_a^b (\partial_\nu w_1)^2(a,\mu)d\nu + \frac{1}{4}\int_a^b (\partial_\mu w_1)^2(\mu,b)d\mu - \frac{1}{2}\int_{\Omega_{ab}} (\mu+\nu)^{-2}|B_2||w_1|^2 d\mu d\nu \leq \\ &a^{-1}q_1^2(a) + b^{-1}q_1^2(b) + \frac{1}{2\sqrt{2}}\int_a^b \left((q_1' + q_2)^2 + q_2^2 + \frac{1}{2}\mu^{-1}B_1(\mu)q_1^2 \right) d\mu. \end{aligned}$$

If we drop the second integral from this inequality and integrate the remaining terms in a , with $a_0 \leq a \leq b$, and use (2.10) we get that

$$\begin{aligned} &\frac{1}{4}\int_{\Omega_{a_0b}} (\partial_\nu w_1)^2 d\nu - \int_{\Omega_{a_0b}} \mu(\mu+\nu)^{-2}|B_2||w_1|^2 d\mu d\nu \leq \\ &\int_{a_0}^b \mu^{-1}q_1^2(\mu)d\mu + q_1^2(b) + \frac{T}{2\sqrt{2}}\int_{a_0}^b ((q_1' + q_2)^2 + q_2^2 + q_1^2) d\mu. \end{aligned}$$

We can use Lemma 2.3 to show that

$$(2.16) \quad \int_{\Omega_{a_0b}} (\mu+\nu)^{-1}|w_1|^2 d\mu d\nu \leq T \int_{\Omega_{a_0b}} (\partial_\nu w_1)^2 d\nu + \int_{a_0}^b \mu^{-1}q_1^2(\mu) d\mu.$$

and therefore, if T_0 is small,

$$(2.17) \quad \frac{1}{8}\int_{\Omega_{a_0b}} (\partial_\nu w_1)^2 d\nu \leq q_1^2(b) + C(\|q_1\|_{1,\frac{1}{2}}^2 + \|q_2\|_{1,\frac{1}{2}}^2).$$

Now we substitute (2.17) and (2.16) into (2.15) and use that $\|q_1\|_{L^\infty} \leq \|q_1\|_{1,0}$ and deduce that

$$\frac{1}{4}\int_a^b (\partial_\nu w_1)^2(a,\nu)d\nu + \frac{1}{4}\int_a^b (\partial_\mu w_1)^2(\mu,b)d\mu \leq C(\|q_1\|_{1,\frac{1}{2}}^2 + \|q_2\|_{1,\frac{1}{2}}^2).$$

By symmetry this estimate also holds in the region below the diagonal. This proves (2.14). \square

Now we prove Proposition 2.6. For $\nu \geq \mu$ we write

$$w_1(\mu, \nu) = w_1(\mu, \mu) + \int_\mu^\nu \partial_s w_1(\mu, s) ds.$$

By the Cauchy-Schwartz inequality,

$$|w_1(\mu, \nu)|^2 \leq 2|w_1(\mu, \mu)|^2 + 2(\nu - \mu) \int_\mu^\nu |\partial_s w_1(\mu, s)|^2 ds,$$

and we deduce from (2.14) that if $(\mu, \nu) \in \Omega_T$, with $T < T_0$ small,

$$(\mu + \nu)^{-1}|w_1(\mu, \nu)|^2 \leq 2\mu^{-1}q_1^2(\mu) + 2 \int_\mu^\nu |\partial_s w_1(\mu, s)|^2 ds \leq C(\|q_1\|_{1,\frac{1}{2}}^2 + \|q_2\|_{1,\frac{1}{2}}^2).$$

By symmetry with respect to the diagonal,

$$|w_1(\mu, \nu)| \leq C(\|q_1\|_{1,\frac{1}{2}}^2 + \|q_2\|_{1,\frac{1}{2}}^2)^{\frac{1}{2}}(\mu + \nu)^{\frac{1}{2}}, \text{ in } \Omega_{T_0}.$$

From now on we will use $C(q_1, q_2)$ to denote a constant which depends on the norms $\|q_1\|_{s,j}$ and $\|q_2\|_{s,j}$ for some s and j . We then go back to equation (2.13) and deduce that if $(\mu, \nu) \in \Omega_{T_0}$,

$$|\partial_\mu \partial_\nu w_1| = |2\mu\nu(\mu + \nu)^{-3} B_1 w_1| \leq C(q_1, q_2)(\mu + \nu)^{-\frac{1}{2}}$$

thus

$$\begin{aligned} |\partial_\nu w_1(\mu, \nu)| &\leq |\partial_\nu w_1(\nu, \nu)| + \int_\mu^\nu |\partial_s \partial_\nu w_1(s, \nu)| ds \leq \\ &|\partial_\nu w_1(\nu, \nu)| + C(q_1, q_2) \int_\mu^\nu (\mu + s)^{-\frac{1}{2}} ds \leq C(q_1, q_2)(\mu + \nu)^{\frac{1}{2}}. \end{aligned}$$

A similar argument shows that

$$|\partial_\mu w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^{\frac{1}{2}}, \text{ if } \nu \geq \mu.$$

By symmetry these estimates hold below the diagonal. This implies that

$$|w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^{\frac{3}{2}} \text{ in } \Omega_{T_0}.$$

We then differentiate equation (2.13) and find that

$$\partial_\mu \partial_\nu^2 w_1 = [\mu(2\mu - \nu)(\mu + \nu)^{-4} B_1 - 2\mu^3 \nu(\mu + \nu)^{-5} B_1'] w_1 + 2\mu\nu(\mu + \nu)^{-1} B_1 \partial_\nu w_1.$$

We deduce that

$$|\partial_\mu \partial_\nu^2 w_1| \leq C(q_1, q_2)(\mu + \nu)^{-\frac{1}{2}}$$

which implies that

$$|\partial_\mu \partial_\nu w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^{\frac{1}{2}}, \quad |\partial_\nu^2 w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^{\frac{1}{2}}, \quad \text{in } \Omega_{T_0}.$$

A similar argument gives that

$$|\partial_\mu^2 w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^{\frac{1}{2}}, \quad \text{in } \Omega_{T_0}.$$

hence we deduce that

$$|\partial_\mu w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^{\frac{3}{2}}, \quad |\partial_\nu w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^{\frac{3}{2}}.$$

This argument can be repeated to show that

$$|\partial_\mu^j \partial_\nu^k w_1(\mu, \nu)| \leq C(q_1, q_2)(\mu + \nu)^m, \quad j, k, m \in \mathbb{N} \text{ in } \Omega_{T_0}.$$

This implies the claim of Proposition 2.6. \square

Next we study the non-degenerate cases, i.e either $n > 3$ or if $n = 3$, $\int_{\partial X} w = 0$, i.e w is orthogonal to the first eigenfunction.

Lemma 2.8. *Let $\Omega_T = (0, T) \times (0, T)$ and let $W \in C^\infty(\Omega_T)$ satisfy*

$$\begin{aligned} (2.18) \quad &\left((\mu + \nu)^2 \partial_\mu \partial_\nu - \phi\left(\frac{2\mu\nu}{\mu + \nu}\right) \Delta_{h_0} - B\left(\frac{2\mu\nu}{\mu + \nu}\right) \right) W = G(\mu, \nu) \text{ in } \Omega_T \times \partial X \\ &W(\mu, \mu) = q_1(\mu), \quad \partial_\mu W(\mu, \mu) = q_2(\mu). \end{aligned}$$

If $n = 3$ we assume that $\int_{\partial X} W = 0$. Then there exists $T_0 > 0$, depending on B and ϕ , and a constant C depending on B and ϕ such that if $\|G\|_{0,2} < \infty$, and $\|q_j\|_{1,1} < \infty$, $j = 1, 2$,

(2.19)

for any fixed $\mu \in [0, T]$

$$\int_0^T \int_{\partial X} (|\partial_\nu W(\mu, \nu)|^2 + (\mu + \nu)^{-2} (|\nabla_{h_0} W(\mu, \nu)|^2 + |W(\mu, \nu)|^2) d\nu \leq C(\|q_1\|_{1,1}^2 + \|q_2\|_{1,1}^2 + \|G\|_{0,2}^2),$$

and for any fixed $\nu \in [0, T]$

$$\int_0^T \int_{\partial X} (|\partial_\nu W(\mu, \nu)|^2 + (\mu + \nu)^{-2} (|\nabla_{h_0} W(\mu, \nu)|^2 + |W(\mu, \nu)|^2) d\mu \leq C(\|q_1\|_{1,1}^2 + \|q_2\|_{1,1}^2 + \|G\|_{0,2}^2).$$

If $\|q_1\|_{1,\frac{3}{2}} < \infty$ and $\|q_2\|_{1,\frac{3}{2}} < \infty$, we also have

$$(2.20) \quad \int_{\Omega_T \times \partial X} (\mu + \nu)^{-2} (\mu |\partial_\mu W|^2 + \nu |\partial_\nu W|^2) + (\mu + \nu)^{-3} (|\nabla_{h_0} W|^2 + |W|^2) d\mu d\nu d\text{vol}_{h_0} \leq CT(\|q_1\|_{2,\frac{3}{2}}^2 + \|q_2\|_{2,\frac{3}{2}}^2 + \|G\|_{0,2}^2).$$

If $\|q_j\|_{2,2} < \infty$, $j = 1, 2$ and $\|G\|_{0,\frac{5}{2}} < \infty$, then

$$(2.21) \quad \int_{\Omega_T \times \partial X} (\mu + \nu)^{-4} (|\nabla_{h_0} W|^2 + |W|^2) d\mu d\nu d\text{vol}_{h_0} \leq C\|G\|_{0,\frac{5}{2}} + CT(\|q_1\|_{2,2}^2 + \|q_2\|_{2,2}^2).$$

Proof. To prove these estimates we multiply (2.18) by $(\mu + \nu)^{-m} (\partial_\mu - \partial_\nu) W$, with $m \in \mathbb{R}_+$, and integrate the product in Ω_{ab} , with $a \leq b \leq T$. The product is equal to

$$\begin{aligned} & \frac{1}{2} \partial_\nu [(\mu + \nu)^{2-m} |\partial_\mu W|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} W|^2 + B |W|^2)] - \\ & \frac{1}{2} \partial_\mu [(\mu + \nu)^{2-m} |\partial_\nu W|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} W|^2 + B |W|^2)] + \\ & \frac{m-2}{2} (\mu + \nu)^{1-m} ((\partial_\mu W)^2 - (\partial_\nu W)^2) + \text{div}_{h_0} [(\mu + \nu)^{-m} \phi \nabla_{h_0} W (\partial_\mu - \partial_\nu) W] + \\ & (\nu - \mu) (\mu + \nu)^{-m-1} (\phi' |\nabla_{h_0} W|^2 + B' |W|^2), \end{aligned}$$

where ϕ' and B' denote the derivative of ϕ and B . Integrating this in $\Omega_{ab} \times \partial X$ and using the divergence theorem and the fact that $(\partial_\nu W)(\mu, \mu) = q'_1(\mu) - q_2(\mu)$,

$$(2.22) \quad \begin{aligned} & \frac{1}{2} \int_{\Sigma_1 \times \partial X} [(\mu + \nu)^{2-m} |\partial_\nu W|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} W|^2 + B |W|^2)] d\nu d\text{vol}_{h_0} - \\ & \frac{1}{2} \int_{\Sigma_2 \times \partial X} [(\mu + \nu)^{2-m} |\partial_\mu W|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} W|^2 + B |W|^2)] d\mu d\text{vol}_{h_0} + \\ & \frac{m-2}{2} \int_{\Omega_{ab} \times \partial X} (\mu + \nu)^{1-m} ((\partial_\mu W)^2 - (\partial_\nu W)^2) d\mu d\nu d\text{vol}_{h_0} + \\ & \int_{\Omega_{ab} \times \partial X} (\nu - \mu) (\mu + \nu)^{-1-m} (\phi' |\nabla_{h_0} W|^2 + B' |W|^2) d\mu d\nu d\text{vol}_{h_0} = \\ & \int_{\Omega_{ab} \times \partial X} G (\mu + \nu)^{-m} (\partial_\mu - \partial_\nu) W d\mu d\nu d\text{vol}_{h_0} + \\ & \frac{1}{2\sqrt{2}} \int_{\Sigma_3 \times \partial X} (2\mu)^{2-m} ((q_2')^2 + (q_2' - q_1)^2) + (2\mu)^{-m} (\phi(\mu) |\nabla_{h_0} q_1|^2 + B(\mu) |q_1|^2) d\mu d\text{vol}_{h_0}. \end{aligned}$$

Here

$$(2.23) \quad \begin{aligned} \Sigma_1 &= \Sigma_1(a, b) = \{(a, \nu), a \leq \nu \leq b\}, \quad \Sigma_2 = \Sigma_2(a, b) = \{(\mu, b), 0 \leq \mu \leq b\} \text{ and} \\ \Sigma_3 &= \Sigma_3(a, b) = \{(\mu, \nu), \mu = \nu, 0 \leq \mu \leq b\}. \end{aligned}$$

When $n > 3$, and T is small, B is positive, but as we saw before, when $n = 3$ this is not necessarily the case. So when $n > 3$ we guarantee that the first two integrals in (2.22) are positive. When $n = 3$, $B(0) = 0$. But we assumed that in this case $\int_{\partial X} W d\text{vol}_{h_0} = 0$ and therefore

$$\int_{\partial X} |\nabla_{h_0} W|^2 d\text{vol}_{h_0} \geq \lambda_2 \int_{\partial X} |W|^2 d\text{vol}_{h_0}.$$

Since $\lambda_2 > 0$, if T_0 is small the term in $B|W|^2$ can be absorbed by the term in $|\nabla_{h_0} W|^2$.

So we may assume that the second integral is positive. We drop it from (2.22) and integrate the remaining terms in the variable a , which determines Σ_1 , with $a_0 \leq a \leq b$. Using (2.10) we obtain

$$(2.24) \quad \begin{aligned} & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} ([(\mu + \nu)^{2-m} - (m-2)\mu(\mu + \nu)^{1-m}] |\partial_\nu W|^2 + (m-2)(\mu + \nu)^{1-m} \mu |\partial_\mu W|^2) d\mu d\nu d\text{vol}_{h_0} + \\ & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-m} (\phi |\nabla_{h_0} W|^2 + B|W|^2) d\mu d\nu d\text{vol}_{h_0} - \\ & \int_{\Omega_{a_0 b} \times \partial X} \mu(\nu - \mu)(\mu + \nu)^{-1-m} (|\phi'| |\nabla_{h_0} W|^2 + |B'| |W|^2) d\mu d\nu d\text{vol}_{h_0} \leq \\ & T(|q_1|_{1, \frac{m}{2}} + |q_2|_{1, \frac{m}{2}}) + \int_{\Omega_{a_0 b} \times \partial X} G\mu(\mu + \nu)^{-m} (\partial_\mu - \partial_\nu) W d\mu d\nu d\text{vol}_{h_0}. \end{aligned}$$

When $m = 2$ this gives

$$\begin{aligned} & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} [|\partial_\nu W|^2 + (\mu + \nu)^{-2} (\phi |\nabla_{h_0} W|^2 + B|W|^2)] d\mu d\nu d\text{vol}_{h_0} - \\ & \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-1} (|\phi'| |\nabla_{h_0} W|^2 + |B'| |W|^2) d\mu d\nu d\text{vol}_{h_0} \leq \\ & T(|q_1|_{1,1} + |q_2|_{1,1}) + T \int_{\Omega_{a_0 b} \times \partial X} |G|(\mu + \nu)^{-2} (\partial_\mu - \partial_\nu) W d\mu d\nu d\text{vol}_{h_0}. \end{aligned}$$

If $n > 3$, then $B > 0$ and if T is small, the second integral can be absorbed into the first. When $n = 3$ the argument used above shows that this can also be done. So we obtain

$$\begin{aligned} & \frac{1}{4} \int_{\Omega_{a_0 b} \times \partial X} [|\partial_\nu W|^2 + (\mu + \nu)^{-2} (\phi |\nabla_{h_0} W|^2 + B|W|^2)] d\mu d\nu d\text{vol}_{h_0} \leq \\ & T(|q_1|_{1,1} + |q_2|_{1,1}) + T \int_{\Omega_{a_0 b} \times \partial X} |G|(\mu + \nu)^{-2} |(\partial_\mu - \partial_\nu) W| d\mu d\nu d\text{vol}_{h_0}. \end{aligned}$$

Now we repeat this argument by dropping the second integral in (2.22) and integrate the remaining terms in b with $a_0 \leq b \leq T$.

$$\begin{aligned} & \frac{1}{4} \int_{\Omega_{a_0 T} \times \partial X} [|\partial_\mu W|^2 + (\mu + \nu)^{-2} (\phi |\nabla_{h_0} W|^2 + B|W|^2)] d\mu d\nu d\text{vol}_{h_0} \leq \\ & T(|q_1|_{1,1} + |q_2|_{1,1}) + T \int_{\Omega_{a_0 T} \times \partial X} |G|(\mu + \nu)^{-2} |(\partial_\mu - \partial_\nu) W| d\mu d\nu d\text{vol}_{h_0}. \end{aligned}$$

We now add these estimates

$$\begin{aligned} & \frac{1}{4} \int_{\Omega_{a_0 T} \times \partial X} [|\partial_\nu W|^2 + |\partial_\mu W|^2 + (\mu + \nu)^{-2}(\phi|\nabla_{h_0} W|^2 + B|W|^2)] d\mu d\nu d\text{vol}_{h_0} \leq \\ & 2T(\|q_1\|_{1,1} + \|q_2\|_{1,1}) + 2T \int_{\Omega_{a_0 T} \times \partial X} |G|(\mu + \nu)^{-2}(\partial_\mu - \partial_\nu)W d\mu d\nu d\text{vol}_{h_0}. \end{aligned}$$

By the Cauchy-Schwartz inequality

$$\int_{\Omega_{a_0 T} \times \partial X} |G|(\mu + \nu)^{-2}(\partial_\mu - \partial_\nu)W d\mu d\nu d\text{vol}_{h_0} \leq 2\|G\|_{0,2} + 4 \int_{\Omega_{a_0 T}} (|\partial_\mu W|^2 + |\partial_\nu W|^2) d\mu d\nu d\text{vol}_{h_0}$$

If T is small this gives

$$\begin{aligned} & \int_{\Omega_{a_0 T} \times \partial X} [|\partial_\nu W|^2 + |\partial_\mu W|^2 + (\mu + \nu)^{-2}(\phi|\nabla_{h_0} W|^2 + B|W|^2)] d\mu d\nu d\text{vol}_{h_0} \leq \\ & CT(\|q_1\|_{1,1} + \|q_2\|_{1,1} + \|G\|_{0,2}). \end{aligned}$$

Equation (2.19) then follows from (2.22).

When $m = 3$ equation (2.24) gives

$$\begin{aligned} & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-2}(\mu|\partial_\mu W|^2 + \nu|\partial_\nu W|^2) d\mu d\nu d\text{vol}_{h_0} + \\ & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-3}(\phi|\nabla_{h_0} W|^2 + B|W|^2) d\mu d\nu d\text{vol}_{h_0} - \\ & \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-2}(|\phi'| |\nabla_{h_0} W|^2 + |B'| |W|^2) d\mu d\nu d\text{vol}_{h_0} \leq \\ & T(\|q_1\|_{1,\frac{3}{2}} + \|q_2\|_{1,\frac{3}{2}}) + \int_{\Omega_{a_0 b} \times \partial X} |G\mu(\mu + \nu)^{-3}(\partial_\mu - \partial_\nu)W| d\mu d\nu d\text{vol}_{h_0}. \end{aligned}$$

The Cauchy-Schwartz inequality gives

$$\begin{aligned} & \int_{\Omega_{a_0 b} \times \partial X} |G\mu(\mu + \nu)^{-3}(\partial_\mu - \partial_\nu)W| d\mu d\nu d\text{vol}_{h_0} \leq 16T\|G\|_{0,2}^2 + \frac{1}{4} \int_{\Omega_{a_0 b}} \mu(\mu + \nu)^{-2}(|\partial_\mu W|^2 + |\partial_\nu W|^2) d\mu d\nu d\text{vol}_{h_0} \leq \\ & 16T\|G\|_{0,2}^2 + \frac{1}{4} \int_{\Omega_{a_0 b}} (\mu + \nu)^{-2}(\mu|\partial_\mu W|^2 + \nu|\partial_\nu W|^2) d\mu d\nu d\text{vol}_{h_0} \leq \end{aligned}$$

Thus we obtain

$$\begin{aligned} & \frac{1}{4} \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-2}(\mu|\partial_\mu W|^2 + \nu|\partial_\nu W|^2) d\mu d\nu d\text{vol}_{h_0} + \\ & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-3}(\phi|\nabla_{h_0} W|^2 + B|W|^2) d\mu d\nu d\text{vol}_{h_0} - \\ & \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-2}(|\phi'| |\nabla_{h_0} W|^2 + |B'| |W|^2) d\mu d\nu d\text{vol}_{h_0} \leq \\ & T(\|q_1\|_{1,\frac{3}{2}} + \|q_2\|_{1,\frac{3}{2}} + \|G\|_{0,2}) \end{aligned}$$

For small T_0 , the third integral can be absorbed into the second and we obtain (2.20).

Now we consider the case when $m = 4$. We deduce from (2.24) that

$$\begin{aligned}
 (2.25) \quad & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} [(\nu + \mu)^{-3}((\nu - \mu)|\partial_\nu W|^2 + 2\mu|\partial_\mu W|^2)] d\mu d\nu d \text{vol}_{h_0} + \\
 & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-4}(\phi|\nabla_{h_0} W|^2 + B|W|^2) d\mu d\nu d \text{vol}_{h_0} - \\
 & \int_{\Omega_{a_0 b} \times \partial X} \mu(\nu - \mu)(\mu + \nu)^{-5}(|\phi'| |\nabla_{h_0} W|^2 + |B'| |W|^2) d\mu d\nu d \text{vol}_{h_0} \leq \\
 & T(\|q_1\|_{1,2} + \|q_2\|_{1,2}) + \int_{\Omega_{a_0 b} \times \partial X} |G| \mu(\mu + \nu)^{-4} |(\partial_\mu - \partial_\nu) W| d\mu d\nu d \text{vol}_{h_0}.
 \end{aligned}$$

We deduce from the Cauchy-Schwartz inequality that

$$\begin{aligned}
 & \int_{\Omega_{a_0 b} \times \partial X} |G| \mu(\mu + \nu)^{-4} |(\partial_\mu - \partial_\nu) W| d\mu d\nu d \text{vol}_{h_0} \leq \\
 & 64 \|G\|_{0, \frac{5}{2}}^2 + \frac{1}{4} \int_{\Omega_{a_0 b}} \mu^2 (\mu + \nu)^{-3} (|\partial_\mu W|^2 + |\partial_\nu W|^2) d\mu d\nu d \text{vol}_{h_0} \leq \\
 & 64 \|G\|_{0, \frac{5}{2}}^2 + \frac{1}{4} \int_{\Omega_{a_0 b}} (\mu + \nu)^{-2} (\mu |\partial_\mu W|^2 + \nu |\partial_\nu W|^2) d\mu d\nu d \text{vol}_{h_0},
 \end{aligned}$$

and from (2.20) we get that

$$\int_{\Omega_{a_0 b} \times \partial X} |G| \mu(\mu + \nu)^{-4} |(\partial_\mu - \partial_\nu) W| d\mu d\nu d \text{vol}_{h_0} \leq C \|G\|_{0, \frac{5}{2}} + CT(\|q_1\|_{1,2} + \|q_2\|_{1,2} + \|G\|_{0,2}).$$

By substituting this into (2.25) we find that

$$\begin{aligned}
 (2.26) \quad & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} [(\nu + \mu)^{-3}((\nu - \mu)|\partial_\nu W|^2 + 2\mu|\partial_\mu W|^2)] d\mu d\nu d \text{vol}_{h_0} + \\
 & \frac{1}{2} \int_{\Omega_{a_0 b} \times \partial X} (\mu + \nu)^{-4}(\phi|\nabla_{h_0} W|^2 + B|W|^2) d\mu d\nu d \text{vol}_{h_0} - \\
 & \int_{\Omega_{a_0 b} \times \partial X} \mu(\nu - \mu)(\mu + \nu)^{-5}(|\phi'| |\nabla_{h_0} W|^2 + |B'| |W|^2) d\mu d\nu d \text{vol}_{h_0} \leq \\
 & T(\|q_1\|_{1,2} + \|q_2\|_{1,2} + \|G\|_{2,0}) + \|G\|_{0, \frac{5}{2}}
 \end{aligned}$$

Again, when T is small the third integral can be absorbed into the second and in particular (2.21) follows from (2.26). This proves Lemma 2.8. \square

Now we are ready to prove Theorem 2.2.

Proof. We will concentrate on the cases not covered by Proposition 2.6. So we assume that if $n = 3$, $\int_{\partial X} w d \text{vol}_{h_0} = 0$.

We can then apply Lemma 2.8 to equation (2.18) with $G = 0$. We get that the solution w to equation (2.5) satisfies

$$(2.27) \quad \|w\|_{0,2} \leq C(\tilde{f}_1, \tilde{f}_2).$$

Since Δ_{h_0} commutes with the equation, we also have

$$(2.28) \quad \|\Delta_{h_0}^k w\|_{0,2} \leq C(\tilde{f}_1, \tilde{f}_2), \quad k = 0, 1, \dots$$

Now we differentiate equation (2.5) with respect to the vector field $\partial_\mu - \partial_\nu$. Let $W_j = (\partial_\mu - \partial_\nu)^j w$. We find that W_1 satisfies

$$\left((\mu + \nu)^2 \partial_\mu \partial_\nu - \phi \left(\frac{2\mu\nu}{\mu + \nu} \right) \Delta_{h_0} - B \left(\frac{2\mu\nu}{\mu + \nu} \right) \right) W_1 = 2(\nu - \mu)(\mu + \nu)^{-1} (\phi' \Delta_{h_0} w - B' w) \text{ in } \Omega_T \times \partial X$$

$$W_1(\mu, \mu) = Q_1(\mu), \quad \partial_\mu W_1(\mu, \mu) = Z_1(\mu),$$

where $Q_1, Z_1 \in C^\infty([0, T])$, depend on \tilde{f}_1 and \tilde{f}_2 and satisfy

$$\partial_\mu^k Q_1(0) = 0, \quad \partial_\mu^k Z_1(0) = 0, \quad k = 0, 1, 2, \dots$$

Let $G_1 = 2(\nu - \mu)(\mu + \nu)^{-1} (\phi' \Delta_{h_0} w - B' w)$. In view of (2.27) and (2.28), $|||G_1|||_{0,2} \leq C(\tilde{f}_1, \tilde{f}_2)$. Then Lemma 2.8 guarantees that $|||W_1|||_{0, \frac{3}{2}} \leq C(\tilde{f}_1, \tilde{f}_2)$. So Lemma 2.5 implies that

$$(2.29) \quad |||w|||_{0, \frac{5}{2}} \leq C(\tilde{f}_1, \tilde{f}_2), \quad |||\Delta_{h_0}^k w|||_{0, \frac{5}{2}} \leq C(\tilde{f}_1, \tilde{f}_2),$$

$$(2.30) \quad k = 0, 1, 2, \dots$$

But then $|||G_1|||_{0, \frac{5}{2}} \leq C(\tilde{f}_1, \tilde{f}_2)$ and Lemma 2.8 guarantees that

$$(2.31) \quad |||W_1|||_{0,2} \leq C(\tilde{f}_1, \tilde{f}_2).$$

Then Lemma 2.5 gives that

$$(2.32) \quad |||w|||_{0,3} \leq C(\tilde{f}_1, \tilde{f}_2), \quad |||\Delta_{h_0} w|||_{0,3} \leq C(\tilde{f}_1, \tilde{f}_2).$$

Now we differentiate (2.5) again with respect to $\partial_\mu - \partial_\nu$. We find that

$$\left((\mu + \nu)^2 \partial_\mu \partial_\nu - \phi \left(\frac{2\mu\nu}{\mu + \nu} \right) \Delta_{h_0} - B \left(\frac{2\mu\nu}{\mu + \nu} \right) \right) W_2 = 2(\nu - \mu)(\mu + \nu)^{-1} (\phi' \Delta_{h_0} - B') W_1 +$$

$$[-4(\mu + \nu)^{-1} (\phi' \Delta_{h_0} - B') + 4(\mu - \nu)^2 (\mu + \nu)^{-2} (\phi'' \Delta_{h_0} - B'')] w \text{ in } \Omega_T \times \partial X$$

$$W_2(\mu, \mu) = Q_2(\mu), \quad \partial_\mu W_2(\mu, \mu) = Z_2(\mu),$$

Let

$$G_2 = 2(\nu - \mu)(\mu + \nu)^{-1} (\phi' \Delta_{h_0} - B') W_1 +$$

$$[-4(\mu + \nu)^{-1} (\phi' \Delta_{h_0} - B') + 4(\mu - \nu)^2 (\mu + \nu)^{-2} (\phi'' \Delta_{h_0} - B'')] w$$

□

In view of (2.31) and (2.32), $|||G_2|||_{0,2} \leq C(\tilde{f}_1, \tilde{f}_2)$, but then it follows from Lemma 2.8 that $|||W_2|||_{0, \frac{3}{2}} \leq C(\tilde{f}_1, \tilde{f}_2)$. Then Lemma 2.5 implies that $|||W_1|||_{0, \frac{5}{2}} \leq C(\tilde{f}_1, \tilde{f}_2)$, and so $|||w|||_{0, \frac{7}{2}} \leq C(\tilde{f}_1, \tilde{f}_2)$. This implies that in fact $|||G_2|||_{0, \frac{5}{2}} \leq C(\tilde{f}_1, \tilde{f}_2)$, and therefore $|||W_2|||_{0,2} \leq C(\tilde{f}_1, \tilde{f}_2)$. Now we differentiate the equation again and repeat the argument. We find that

$$|||(\partial_\mu - \partial_\nu)^j w|||_{0, \frac{5}{2}} \leq C(\tilde{f}_1, \tilde{f}_2), \quad j = 1, 2, \dots$$

and by Lemma 2.5 we conclude that

$$(2.33) \quad |||w|||_{0,j} \leq C(\tilde{f}_1, \tilde{f}_2), \quad |||\Delta_{h_0}^k w|||_{0,j} \leq C(\tilde{f}_1, \tilde{f}_2), \quad k, j = 1, 2, \dots$$

Now we go back to equation (2.22) and apply it to the solution w to (2.5). In this case $G = 0$ and we obtain

$$\begin{aligned}
(2.34) \quad & \frac{1}{2} \int_{\Sigma_1 \times \partial X} [(\mu + \nu)^{2-m} |\partial_\nu w|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} w|^2 + B |w|^2)] d\nu d \text{vol}_{h_0} + \\
& \frac{1}{2} \int_{\Sigma_2 \times \partial X} [(\mu + \nu)^{2-m} |\partial_\mu w|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} w|^2 + B |W|^2)] d\mu d \text{vol}_{h_0} + \\
& \frac{m-2}{2} \int_{\Omega_{ab} \times \partial X} (\mu + \nu)^{1-m} ((\partial_\mu w)^2 - (\partial_\nu w)^2) d\mu d\nu d \text{vol}_{h_0} + \\
& \int_{\Omega_{ab} \times \partial X} (\nu - \mu) (\mu + \nu)^{-1-m} (\phi' |\nabla_{h_0} w|^2 + B' |w|^2) d\mu d\nu d \text{vol}_{h_0} = \\
& \frac{1}{2\sqrt{2}} \int_{\Sigma_3 \times \partial X} (2\mu)^{2-m} ((\tilde{f}_2)^2 + (\tilde{f}_2 - \tilde{f}_1)^2) + (2\mu)^{-m} (\phi(\mu) |\nabla_{h_0} \tilde{f}_1|^2 + B(\mu) |\tilde{f}_1|^2) d\mu d \text{vol}_{h_0}.
\end{aligned}$$

The term

$$\begin{aligned}
& \int_{\Omega_{ab} \times \partial X} (\mu + \nu)^{1-m} ((\partial_\mu w)^2 - (\partial_\nu w)^2) d\mu d\nu d \text{vol}_{h_0} = \int_{\Omega_{ab} \times \partial X} (\mu + \nu)^{1-m} ((\partial_\mu - \partial_\nu)w)((\partial_\mu + \partial_\nu)w) d\mu d\nu d \text{vol}_{h_0} \leq \\
& 2 \int_{\Omega_{ab} \times \partial X} (\mu + \nu)^{2-2m} |(\partial_\mu - \partial_\nu)w|^2 d\mu d\nu d \text{vol}_{h_0} + 2 \int_{\Omega_{ab} \times \partial X} |(\partial_\mu + \partial_\nu)w|^2 d\mu d\nu d \text{vol}_{h_0} \leq C(\tilde{f}_1, \tilde{f}_2),
\end{aligned}$$

and from (2.33) we have that

$$\left| \int_{\Omega_{ab} \times \partial X} (\nu - \mu) (\mu + \nu)^{-1-m} (\phi' |\nabla_{h_0} w|^2 + B' |w|^2) d\mu d\nu d \text{vol}_{h_0} \right| \leq C(\tilde{f}_1, \tilde{f}_2).$$

Therefore we conclude that

$$\begin{aligned}
(2.35) \quad & \frac{1}{2} \int_{\Sigma_1 \times \partial X} [(\mu + \nu)^{2-m} |\partial_\nu w|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} w|^2 + B |w|^2)] d\nu d \text{vol}_{h_0} + \\
& \frac{1}{2} \int_{\Sigma_2 \times \partial X} [(\mu + \nu)^{2-m} |\partial_\mu w|^2 + (\mu + \nu)^{-m} (\phi |\nabla_{h_0} w|^2 + B |W|^2)] d\mu d \text{vol}_{h_0} \leq C(\tilde{f}_1, \tilde{f}_2).
\end{aligned}$$

Now we write, for $m > 2$,

$$\begin{aligned}
(2.36) \quad & |(\mu + \nu)^{1-m} w(\mu, \nu)| \leq 2\mu^{1-m} |w(\mu, \mu)| + 2(\mu + \nu)^{2-m} \int_\mu^\nu |\partial_s w(\mu, s)|^2 ds \leq \\
& 2\mu^{1-m} |w(\mu, \mu)| + 2 \int_\mu^\nu (\mu + s)^{2-m} |\partial_s w(\mu, s)|^2 ds \leq C(\tilde{f}_1, \tilde{f}_2).
\end{aligned}$$

Now we argue as we did in the case of w_0 , and conclude that $w \in C^\infty(\overline{\Omega_T} \times \partial X)$. This ends the proof of Theorem 2.2. \square

To prove Theorem 1.3 we need the following:

Lemma 2.9. *Suppose that $f \in \mathcal{S}(X)$ and $\mathcal{R}_+(0, f)(s, y) = 0$ for $s < s_0 \ll 0$. Let u be the solution to (2.1) and let w be the function defined by u in (2.4). Then*

$$\begin{aligned}
(2.37) \quad & \partial_\mu^k w(0, \nu, y) = 0 \text{ if } \nu \leq -\frac{1}{s_0}, \quad k = 0, 1, \dots \\
& \partial_\nu^k w(\mu, 0, y) = 0 \text{ if } \mu \leq -\frac{1}{s_0}, \quad k = 0, 1, \dots
\end{aligned}$$

Proof. Suppose $\mathcal{R}_+(0, f)(s, y) = 0$ for $s < s_0$. Since the initial data is of the form $(0, f)$, the solution u of (2.1) is odd in time, and therefore $\mathcal{R}_-(0, f)(s, y) = 0$ if $s > -s_0$. Since w is smooth up to $\{\mu = 0\}$ and $\{\nu = 0\}$, this means that

$$\partial_\nu w(0, \nu, y) = 0 \quad \text{if } \nu \leq -\frac{1}{s_0}, \quad \partial_\mu w(\mu, 0, y) = 0 \quad \text{if } \mu \leq -\frac{1}{s_0}.$$

We know from (2.36) that w vanishes to infinite order at $\{\mu = \nu = 0\}$. Therefore

$$w(0, \nu, y) = 0 \quad \text{if } \nu \leq -\frac{1}{s_0}, \quad w(\mu, 0, y) = 0 \quad \text{if } \mu \leq -\frac{1}{s_0}.$$

From the equation (2.5), we conclude that

$$\begin{aligned} \text{if } \mu \neq 0, \quad \partial_\mu \partial_\nu w(\mu, 0, y) &= \mu^{-2}(\phi(0)\Delta_{h_0} + B(0))w(\mu, 0) = 0 \\ \text{if } \nu \neq 0, \quad \partial_\mu \partial_\nu w(0, \nu, y) &= \nu^{-2}(\phi(0)\Delta_{h_0} + B(0))w(0, \nu) = 0. \end{aligned}$$

Using that w vanishes to infinite order at $\{\mu = \nu = 0\}$,

$$\begin{aligned} \partial_\mu w(0, \nu, y) &= 0 \quad \text{if } \nu \leq -\frac{1}{s_0}, \\ \partial_\nu w(\mu, 0, y) &= 0 \quad \text{if } \mu \leq -\frac{1}{s_0}. \end{aligned}$$

Now we differentiate equation (2.5) with respect to μ we find that

$$(\mu + \nu)^2 \partial_\mu^2 \partial_\nu w + \phi \Delta_{h_0} \partial_\mu w + B \partial_\mu w + 2(\mu + \nu) \partial_\mu \partial_\nu w + \nu^2 (\mu + \nu)^{-2} \Delta_{h_0} w + \nu^2 (\mu + \nu)^{-2} B' w = 0$$

Hence,

$$\text{if } \nu \neq 0, \quad \partial_\mu^2 \partial_\nu w(0, \nu, y) = 0.$$

Since w is smooth, and vanishes to infinite order at $\mu = \nu = 0$,

$$\partial_\mu^2 w(0, \nu, y) = 0 \quad \text{if } \nu \leq -\frac{1}{s_0}.$$

By symmetry,

$$\partial_\nu^2 w(\mu, 0, y) = 0 \quad \text{if } \mu \leq -\frac{1}{s_0}.$$

Since away from $\{\mu = \nu = 0\}$ the coefficients of (2.5) are smooth, we can repeat the argument to prove that all derivatives of w vanish at $\{\mu = 0, \nu \leq -\frac{1}{s_0}\} \cup \{\nu = 0, \mu \leq -\frac{1}{s_0}\}$. \square

3. CARLEMAN ESTIMATES

Let w be a solution to (2.5) with $\tilde{f}_1 = 0$ and $\tilde{f}_2 = \tilde{f} = \frac{1}{2\mu^2} F(\mu) f(\mu, y)$, and let

$$w_k(\mu, \nu) = \langle w(\mu, \nu, y), \phi_k(y) \rangle_{L^2(\partial X, d\text{vol}_{h_0})},$$

where $\phi_k, k = 1, 2, \dots$ are the eigenfunctions of Δ_{h_0} with eigenvalue λ_k , where $0 = \lambda_1 < \lambda_2$, and $\lambda_k \leq \lambda_{k+1}$, for $k > 1$, and $\lambda_k \rightarrow \infty$. Then $w_k \in C^\infty([0, T] \times [0, T])$ satisfies

$$\begin{aligned} ((\mu + \nu)^2 \partial_\mu \partial_\nu + F_k(\mu, \nu)) w_k &= 0 \text{ in } [0, T] \times [0, T], \\ w_k(\mu, \mu) = 0, \quad \partial_\mu w_k(\mu, \mu) &= \tilde{f}_k(\mu) = \langle \tilde{f}, \phi_k \rangle, \quad j = 1, 2. \end{aligned}$$

Here $F_k(\mu, \nu) = \lambda_k \phi(\mu, \nu) + B(\mu, \nu)$.

By assumption w vanishes to infinite order at $\{\mu = 0\}$ and $\{\nu = 0\}$. Thus so does $w_k(\mu, \nu)$, $k = 1, 2, \dots$. We will prove that under these assumptions there exists $\epsilon > 0$, independent of k such that $w_k(\mu, \nu) = 0$ if $\mu \leq \epsilon$ and $\nu \leq \epsilon$. In particular $\tilde{f}_k(\mu) = 0$, $k = 0, 1, 2, \dots$ if $\mu \leq \epsilon$, and hence $f(x) = 0$ if $x \leq \epsilon$.

It is convenient to work with coordinates r and τ defined in (2.12). The main ingredient of the proof is

Lemma 3.1. *Let U be a neighborhood of $(0, 0)$. Let*

$$P_k = r^2 \partial_r^2 - r^2 \partial_\tau^2 + F_k(\tau, r),$$

where $F_k(r, \tau) = \lambda_k \phi(\frac{r^2 - \tau^2}{2r}) - B(\frac{r^2 - \tau^2}{2r})$. Then there exists $C > 0$, independent of k , and $\gamma_0 = \gamma_0(k)$ such that for every $\gamma > \gamma_0$, and every $u \in C_0^\infty(U)$, which is supported in $\{(\tau, r) : r \geq 0 \text{ and } -r \leq \tau \leq r\}$,

$$(3.1) \quad \|r^{-\gamma-2} P_k u\|^2 \geq C (\gamma^2 \|r^{-1} \partial_r r^{-\gamma} u\|^2 + \gamma^2 \|r^{-1-\gamma} \partial_\tau u\|^2 + \gamma^4 \|r^{-\gamma-2} u\|^2).$$

Here $\|\cdot\|$ denotes the $L^2(U)$ norm.

Proof. Let

$$P_{\gamma,k} = r^{-\gamma-2} P_k r^\gamma, \quad P_{\gamma,k} = r^{-2} P_k + 2\gamma r^{-1} \partial_r + \gamma(\gamma-1) r^{-2}.$$

The support of u is contained in $\{r \geq 0\}$ and $\{r \geq \tau \geq -r\}$. So we write

$$u = r^\gamma v, \text{ and } r^{-\gamma-2} P_k u = P_{\gamma,k} v.$$

We have

$$(3.2) \quad \|P_{\gamma,k} v\|^2 = \|r^{-2} P_k v\|^2 + 4\gamma^2 \|r^{-1} \partial_r v\|^2 + \gamma^2 (\gamma-1)^2 \|r^{-2} v\|^2 + 4\gamma \langle r^{-2} P_k v, r^{-1} \partial_r v \rangle + 2\gamma(\gamma-1) \langle r^{-2} P_k v, r^{-2} v \rangle + 4\gamma^2 (\gamma-1) \langle r^{-1} \partial_r v, r^{-2} v \rangle.$$

Now we integrate by parts to compute the inner products $\langle r^{-2} P_k v, r^{-1} \partial_r v \rangle$, $\langle r^{-2} P_k v, r^{-2} v \rangle$ and $\langle r^{-1} \partial_r v, r^{-2} v \rangle$. We begin with

$$\langle r^{-1} \partial_r v, r^{-2} v \rangle = \frac{1}{2} \int r^{-3} \partial_r v^2 dr d\tau = \frac{3}{2} \|r^{-2} v\|^2.$$

$\langle r^{-2} P_k v, r^{-1} \partial_r v \rangle$ has two terms:

$$\langle (\partial_r^2 - \partial_t^2) v, r^{-1} \partial_r v \rangle = \frac{1}{2} \int r^{-1} \partial_r (\partial_r v)^2 dr d\tau - \int r^{-1} \partial_t^2 \partial_r v dr d\tau = \frac{1}{2} \|r^{-1} \partial_r v\|^2 + \frac{1}{2} \|r^{-1} \partial_t v\|^2,$$

and

$$\langle r^{-2} F_k v, r^{-1} \partial_r v \rangle = \frac{1}{2} \int r^{-3} F_k \partial_r v^2 dr d\tau = \frac{3}{2} \int r^{-4} (F_k + r \partial_r F_k) v^2 dr d\tau \geq -C(\lambda_k + 1) \|r^{-2} v\|^2.$$

$\langle r^{-2} P_k v, r^{-1} \partial_r v \rangle$ also has two terms:

$$\langle (\partial_r^2 - \partial_t^2) v, r^{-2} v \rangle = 3 \|r^{-2} v\|^2 - \|r^{-1} \partial_r v\|^2 + \|r^{-1} \partial_t v\|^2.$$

and

$$\langle r^{-2} F_k v, r^{-2} v \rangle \geq -C(\lambda_k + 1) \|r^{-2} v\|^2.$$

Putting these estimates together we find that

$$\begin{aligned} \|P_{\gamma,k} v\|^2 &\geq (2\gamma^2 + 4\gamma) \|r^{-1} \partial_r v\|^2 + 2\gamma^2 \|r^{-1} \partial_t v\|^2 + \\ &(\gamma^2 (\gamma-1)^2 + 6\gamma^2 (\gamma-1) + 6\gamma (\gamma-1) - 4C(\lambda_k + 1)\gamma - 2\gamma(\gamma-1)\lambda_k) \|r^{-2} v\|^2. \end{aligned}$$

Thus, if $\gamma_0 \gg C(\lambda_k + 1)$,

$$\|P_{\gamma,k} v\|^2 \geq C(\gamma^2 \|r^{-1} \partial_r v\|^2 + \gamma^2 \|r^{-1} \partial_t v\|^2 + \gamma^4 \|r^{-2} v\|^2).$$

Since $v = r^{-\gamma} u$ this implies that

$$(3.3) \quad \|r^{-\gamma-2} P_k u\|^2 \geq C(\gamma^2 \|r^{-\gamma} \partial_r r^{-1} u\|^2 + \gamma^2 \|r^{-\gamma-1} \partial_t u\|^2 + \gamma^4 \|r^{-\gamma-2} u\|^2, \quad \gamma > \gamma_0.$$

4. PROOF OF THEOREM 1.3

To prove Theorem 1.3 we apply Lemma 3.1 to $\chi(r)w_k$, with $\chi \in C^\infty(-T, T)$, such that $\chi(r) = 1$ if $|r| < \frac{T}{4}$ and $\chi(r) = 0$ if $|r| > \frac{T}{2}$.

As a consequence of (3.3),

$$\|r^{-\gamma-2}P_k\chi w_k\| \geq C\gamma^4\|r^{-\gamma-2}\chi w_k\|^2.$$

Notice that, since $P_k w_k = 0$,

$$P_k\chi(r)w_k = r^2\chi''(r)w_k + 2r^2\chi'(r)\partial_r w_k,$$

and therefore $P_k\chi(r)w_k$ is supported in $\frac{T}{4} \leq r \leq \frac{T}{2}$. Thus

$$\|r^{-\gamma-2}P_k\chi w_k\|^2 \leq C(\chi, w_k)T^{-2\gamma-2}4^{2\gamma+2}$$

But since $\chi(r) = 1$ if $r < T/4$,

$$\|r^{-\gamma-2}\chi w_k\| \geq \|r^{-\gamma-2}w_k\|_{L^2(B(0, \frac{T}{4}))} \geq T^{-2\gamma-2}4^{2\gamma+2}\|w_k\|_{L^2(B(0, \frac{T}{4}))},$$

where $\|w_k\|_{L^2(B(0, \frac{T}{4}))}$ is the L^2 norm of w_k on the ball centered at $(0, 0)$ and radius $\frac{T}{4}$. So we have

$$C(\chi, w_k) \geq C\gamma^4\|w_k\|_{L^2(B(0, \frac{T}{4}))}, \quad \gamma \geq \gamma_0.$$

Letting $\gamma \rightarrow \infty$ gives

$$\|w_k\|_{L^2(B(0, \frac{T}{4}))} = 0.$$

Since T does not depend on k we conclude that $w = 0$ in $B(0, \frac{T}{4})$. In particular this shows that $f(x) = 0$ if $x \leq \frac{T}{4}$. Therefore f is compactly supported. Then Theorem 1.3 follows from Theorem 1.2.

Notice that the weight in the Carleman estimate depends on the eigenvalue λ_k . The bigger the eigenvalue, the larger the parameter γ has to be.

□

5. ACKNOWLEDGEMENTS

This research was partially funded by the NSF grant DMS 0500788.

This paper was written in the summer 2007 when I visited the Mathematics Department of the Federal University of Pernambuco, Recife, Brazil. I would like to thank Prof. Fernando Cardoso for the hospitality. I would also like to thank the CNPq (Brazil) for the financial support during my stay in Recife.

□

REFERENCES

- [1] F.G. Friedlander, *On the radiation field of pulse solutions of the wave equation*. Proc. Roy. Soc. Ser. A. **269**, 53-65, (1962)
- [2] F.G. Friedlander, *On the radiation field of pulse solutions of the wave equation II*. Proc. Roy. Soc. Ser. A. **279**, 386-394, (1963)
- [3] F.G. Friedlander, *On the radiation field of pulse solutions of the wave equation III*. Proc. Roy. Soc. Ser. A. **299**, 386-394, (1965)
- [4] F.G. Friedlander, *Radiation fields and hyperbolic scattering theory*. Math. Proc. Camb. Phil. Soc., **88**, 483-515, (1980)
- [5] F.G. Friedlander, *Notes on the wave equation on asymptotically Euclidean manifolds*. J. of Func. Anal. **184**, no. 1, 1-18, (2001)
- [6] S. Helgason. *The Radon Transform*. Birkhauser, 2nd edition, (1999).
- [7] L. Hörmander, *The analysis of linear partial differential operators*. Vol 1-4, Grundlehren Math. Wiss. **256**, Springer Verlag, (1983)

- [8] M.S. Joshi and A. Sá Barreto. *Recovering asymptotics of metrics from fixed energy scattering data*. Invent. Math. **137**, 127-143, (1999)
- [9] P. Lax and R. Phillips. *Scattering theory*. Academic Press, 1989, Revised edition .
- [10] R. Mazzeo and R.B. Melrose. *Meromorphic extension of the resolvent on complete spaces with asymptotically constant negative curvature*. J. Func. Anal. **108**, 260-310, (1987)
- [11] R.B. Melrose, *Geometric scattering theory*. Stanford Lectures, Cambridge Univ. Press, (1995)
- [12] R.B. Melrose, *Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces*. Spectral and scattering theory (Sanda, 1992), 85–130, Lecture Notes in Pure and Appl. Math., 161
- [13] A. Sá Barreto, *Radiation fields on asymptotically Euclidean manifolds*. Comm. in P.D.E. **28** , Nos. 9 & 10, 1661-1673, (2003)
- [14] A. Sá Barreto, *Radiation fields and inverse scattering on asymptotically Euclidean manifolds*. Partial differential equations and inverse problems, Contemp. Math., 362, Amer. Math. Soc., Providence, RI, 371–380, 2004.
- [15] A. Sá Barreto, *Radiation fields, scattering, and inverse scattering on asymptotically hyperbolic manifolds*. Duke Math. J. 129, and no. 3, 407–480, (2005)

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 NORTH UNIVERSITY STREET, WEST LAFAYETTE IN 47907, USA

E-mail address: `sabarre@math.purdue.edu`